

CODATA RECOMMENDED VALUES OF THE FUNDAMENTAL PHYSICAL CONSTANTS: 2014

NIST SP 961 (Sept/2015) Values from: P. J. Mohr, D. B. Newell, and B. N. Taylor, arXiv:1507.07956

A more extensive listing of constants is available in the above reference and on the NIST Physics Laboratory Web site physics.nist.gov/constants.

The number in parentheses is the one-standard-deviation uncertainty in the last two digits of the given value.

Quantity	Symbol	Numerical value	Unit	Quantity	Symbol	Numerical value	Unit
speed of light in vacuum	c	299 792 458 (exact)	$m s^{-1}$	muon g-factor $-2(1 + a_\mu)$	g_μ	$-2.002\ 331\ 8418(13)$	
magnetic constant	μ_0	$4\pi \times 10^{-7}$ (exact)	$N A^{-2}$	muon-proton magnetic moment ratio	μ_μ/μ_p	$-3.183\ 345\ 142(71)$	
electric constant $1/\epsilon_0 c^2$	ϵ_0	$= 12.566\ 370\ 614... \times 10^{-7}$	$N A^{-2}$	proton mass	m_p	$1.672\ 621\ 898(21) \times 10^{-27}$	kg
Newtonian constant of gravitation	G	$8.854\ 187\ 817... \times 10^{-12}$	$F m^{-1}$	in u		$1.007\ 276\ 466\ 879(91)$	u
Planck constant	h	$6.626\ 070\ 040(81) \times 10^{-34}$	$J s$	energy equivalent in MeV	$m_p c^2$	$938.272\ 0813(58)$	MeV
in eV s		$4.135\ 667\ 662(25) \times 10^{-15}$	$eV s$	proton-electron mass ratio	m_p/m_e	$1836.152\ 673\ 89(17)$	
$h/2\pi$	\hbar	$1.054\ 571\ 800(13) \times 10^{-34}$	$J s$	proton magnetic moment	μ_p	$1.410\ 606\ 7873(97) \times 10^{-26}$	$J T^{-1}$
in eV s		$6.582\ 119\ 514(40) \times 10^{-16}$	$eV s$	to nuclear magneton ratio	μ_p/μ_N	$2.792\ 847\ 3508(85)$	
elementary charge	e	$1.602\ 176\ 6208(98) \times 10^{-19}$	C	proton magnetic shielding correction $1 - \mu_p/\mu_N$	σ_p	$25.691(11) \times 10^{-6}$	
magnetic flux quantum $h/2e$	Φ_0	$2.067\ 833\ 831(13) \times 10^{-15}$	Wb	(H_2O , sphere, 25 °C)			
Josephson constant $2e/h$	K_J	$483\ 597.8525(30) \times 10^9$	$Hz V^{-1}$	proton gyromagnetic ratio $2\mu_p/\hbar$	γ_p	$2.675\ 221\ 900(18) \times 10^8$	$s^{-1} T^{-1}$
von Klitzing constant $h/e^2 = \mu_0 c/2\alpha$	R_K	$25\ 812.807\ 4555(59)$	Ω	shielded proton gyromagnetic ratio $2\mu_p/\hbar$	γ_p'	$42.577\ 478\ 92(29)$	$MHz T^{-1}$
Bohr magneton $eh/2m_e$	μ_B	$927.400\ 9994(57) \times 10^{-26}$	$J T^{-1}$	(H_2O , sphere, 25 °C)	γ_p'	$2.675\ 153\ 171(33) \times 10^8$	$s^{-1} T^{-1}$
in $eV T^{-1}$		$5.788\ 381\ 8012(26) \times 10^{-5}$	$eV T^{-1}$	neutron mass in u	m_n	$1.008\ 664\ 915\ 88(49)$	u
nuclear magneton $eh/2m_p$	μ_N	$5.050\ 783\ 699(31) \times 10^{-27}$	$J T^{-1}$	energy equivalent in MeV	$m_n c^2$	$939.565\ 4133(58)$	MeV
in $eV T^{-1}$		$3.152\ 451\ 2550(15) \times 10^{-8}$	$eV T^{-1}$	neutron-proton mass ratio	m_n/m_p	$1.001\ 378\ 418\ 98(51)$	
fine-structure constant $e^2/4\pi\epsilon_0\hbar c$	α	$7.297\ 352\ 5664(17) \times 10^{-3}$		neutron magnetic moment	μ_n	$-0.966\ 236\ 50(23) \times 10^{-26}$	$J T^{-1}$
inverse fine-structure constant	α^{-1}	$137.035\ 999\ 139(31)$		to nuclear magneton ratio	μ_n/μ_N	$-1.913\ 042\ 73(45)$	
Rydberg constant $\alpha^2 m_e c^2/2h$	R_∞	$10973\ 731.568\ 508(65)$	m^{-1}	deuteron mass in u	m_d	$2.013\ 553\ 212\ 745(40)$	u
energy equivalent in eV	$R_\infty c$	$3.289\ 841\ 960\ 355(19) \times 10^{15}$	Hz	energy equivalent in MeV	$m_d c^2$	$1875.612\ 928(12)$	MeV
Bohr radius $a_0/4\pi R_\infty = 4\pi\epsilon_0\hbar^2/m_e e^2$	a_0	$0.529\ 177\ 210\ 67(12) \times 10^{-10}$	m	deuteron-proton mass ratio	m_d/m_p	$1.999\ 007\ 500\ 87(19)$	
Hartree energy $e^2/4\pi\epsilon_0 a_0 = 2R_\infty hc = \alpha^2 m_e c^2$	E_h	$4.359\ 744\ 650(54) \times 10^{-18}$	J	deuteron magnetic moment	μ_d	$0.433\ 073\ 5040(36) \times 10^{-26}$	$J T^{-1}$
in eV		$27.211\ 386\ 02(17)$	eV	to nuclear magneton ratio	μ_d/μ_N	$0.857\ 438\ 2311(48)$	
electron mass	m_e	$9.109\ 383\ 56(11) \times 10^{-31}$	kg	helium (3He nucleus) mass in u	m_3	$3.014\ 932\ 246\ 73(12)$	u
in u		$5.485\ 799\ 090\ 70(16) \times 10^{-4}$	u	energy equivalent in MeV	$m_3 c^2$	$2808.391\ 586(17)$	MeV
energy equivalent in MeV	$m_e c^2$	$0.510\ 998\ 9461(31)$	MeV	shielded helium magnetic moment (gas, sphere, 25 °C)	μ_3	$-1.074\ 553\ 080(14) \times 10^{-26}$	$J T^{-1}$
electron-muon mass ratio	m_e/m_μ	$4.836\ 331\ 70(11) \times 10^{-3}$		to Bohr magneton ratio	μ_3/μ_B	$-1.158\ 671\ 471(14) \times 10^{-3}$	
electron-proton mass ratio	m_e/m_p	$5.446\ 170\ 213\ 52(52) \times 10^{-4}$		to nuclear magneton ratio	μ_3/μ_N	$-2.127\ 497\ 720(25)$	
electron charge to mass quotient	$-e/m_e$	$-1.758\ 820\ 024(11) \times 10^{11}$	$C kg^{-1}$	alpha particle mass in u	m_α	$4.001\ 506\ 179\ 127(63)$	u
Compton wavelength $h/m_e c$	λ_C	$2.426\ 310\ 2367(11) \times 10^{-12}$	m	energy equivalent in MeV	$m_\alpha c^2$	$3727.379\ 378(23)$	MeV
$\lambda_C/2\pi = \alpha a_0 = \alpha^2/4\pi R_\infty$	λ_C	$386.159\ 267\ 64(18) \times 10^{-15}$	m	Avogadro constant	N_A, L	$6.022\ 140\ 857(74) \times 10^{23}$	mol^{-1}
classical electron radius $\alpha^2 a_0$	r_e	$2.817\ 940\ 3227(19) \times 10^{-15}$	m	atomic mass constant $\frac{1}{12}m(^{12}C) = 1 u$	m_u	$1.660\ 539\ 040(20) \times 10^{-27}$	kg
Thomson cross section $(8\pi/3)r_e^2$	σ_e	$0.665\ 245\ 871\ 58(91) \times 10^{-28}$	m^2	energy equivalent in MeV	$m_u c^2$	$931.494\ 0954(57)$	MeV
electron magnetic moment	μ_e	$-928.476\ 4620(57) \times 10^{-26}$	$J T^{-1}$	Faraday constant $N_A e$	F	$96\ 485.332\ 89(59)$	$C mol^{-1}$
to Bohr magneton ratio	μ_e/μ_B	$-1.001\ 159\ 652\ 180\ 91(26)$		R	$8.314\ 4598(48)$	$J K^{-1} mol^{-1}$	
to nuclear magneton ratio	μ_e/μ_N	$-1838.281\ 972\ 34(17)$		Boltzmann constant R/N_A	k	$1.380\ 648\ 52(79) \times 10^{-23}$	$J K^{-1}$
electron magnetic moment anomaly $ \mu_e /\mu_B - 1$	a_e	$1.159\ 652\ 180\ 91(26) \times 10^{-3}$		in $eV K^{-1}$		$8.617\ 3303(50) \times 10^{-5}$	$eV K^{-1}$
electron g-factor $-2(1 + a_e)$	g_e	$-2.002\ 319\ 304\ 361\ 82(52)$		molar volume of ideal gas RT/p	V_m	$22.413\ 962(13) \times 10^{-3}$	$m^3 mol^{-1}$
electron-proton magnetic moment ratio	μ_e/μ_p	$-658.210\ 6866(20)$		($T = 273.15 K, p = 101.325 kPa$)			
muon mass in u	m_μ	$0.113\ 428\ 9257(25)$	u	Stefan-Boltzmann constant $\pi^2 k^4/60\hbar^3 c^2$	σ	$5.670\ 367(13) \times 10^{-8}$	$W m^{-2} K^{-4}$
energy equivalent in MeV	$m_\mu c^2$	$105.658\ 3745(24)$	MeV	first radiation constant $2\pi^5 hc^2/15$	c_1	$3.741\ 771\ 790(46) \times 10^{-16}$	$W m^2$
muon-electron mass ratio	m_μ/m_e	$206.768\ 2826(46)$		second radiation constant hc/k	c_2	$1.438\ 777\ 36(83) \times 10^{-2}$	$m K$
muon magnetic moment	μ_μ	$-4.490\ 448\ 26(10) \times 10^{-26}$	$J T^{-1}$	Wien displacement law constant $b = \lambda_{max} T = c_2/4.965\ 114\ 231...$		$2.897\ 7729(17) \times 10^{-3}$	$m K$
to Bohr magneton ratio	μ_μ/μ_B	$-4.841\ 970\ 48(11) \times 10^{-3}$		Cu x unit: $\lambda(Cu K\alpha_1)/1537.400$	$xu(Cu K\alpha_1)$	$1.002\ 076\ 97(28) \times 10^{-13}$	m
to nuclear magneton ratio	μ_μ/μ_N	$-8.890\ 597\ 05(20)$		Mo x unit: $\lambda(Mo K\alpha_1)/707.831$	$xu(Mo K\alpha_1)$	$1.002\ 099\ 52(53) \times 10^{-13}$	m
muon magnetic moment anomaly $ \mu_\mu /(\epsilon\hbar/2m_\mu) - 1$	a_μ	$1.165\ 920\ 89(63) \times 10^{-3}$					
Energy equivalents							
$(1 m^{-1})c = 299\ 792\ 458 Hz$	$(1 Hz)h/k = 4.799\ 2447(28) \times 10^{-11} K$	$(1 J) = 6.241\ 509\ 126(38) \times 10^{18} eV$	$(1 eV)/c^2 = 1.073\ 544\ 1105(66) \times 10^{-9} u$				
$(1 m^{-1})hc/k = 1.438\ 777\ 36(83) \times 10^{-2} K$	$(1 Hz)h = 4.135\ 667\ 662(25) \times 10^{-15} eV$	$(1 eV) = 1.602\ 176\ 6208(98) \times 10^{-19} J$	$(1 kg) = 6.022\ 140\ 857(74) \times 10^{26} u$				
$(1 m^{-1})hc = 1.239\ 841\ 9739(76) \times 10^{-6} eV$	$(1 K)k/hc = 69.503\ 457(40) m^{-1}$	$(1 eV)/hc = 8.065\ 544\ 005(50) \times 10^5 m^{-1}$	$(1 u) = 1.660\ 539\ 040(20) \times 10^{-27} kg$				
$(1 m^{-1})h/c = 1.331\ 025\ 049\ 00(61) \times 10^{-15} u$	$(1 K)k/h = 2.083\ 6612(12) \times 10^{10} Hz$	$(1 eV)/h = 2.417\ 989\ 262(15) \times 10^{14} Hz$	$(1 u)/c\hbar = 7.513\ 006\ 6166(34) \times 10^{14} m^{-1}$				
$(1 Hz)/c = 3.335\ 640\ 951... \times 10^{-9} m^{-1}$	$(1 K)k = 8.617\ 3303(50) \times 10^{-5} eV$	$(1 eV)/k = 1.160\ 452\ 21(67) \times 10^4 K$	$(1 u)c^2 = 931.494\ 0954(57) \times 10^6 eV$				